
**UNITED STATES
SECURITIES AND EXCHANGE COMMISSION**

Washington, D.C. 20549

FORM 8-K

**CURRENT REPORT
Pursuant to Section 13 or 15(d)
of the Securities Exchange Act of 1934**

Date of Report (Date of earliest event reported): September 16, 2017

PIERIS PHARMACEUTICALS, INC.

(Exact Name of Registrant as Specified in its Charter)

Nevada
(State of Incorporation)

001-37471
(Commission
File Number)

EIN 30-0784346
(IRS Employer
Identification No.)

**255 State Street, 9th Floor
Boston, MA
United States**
(Address of principal executive offices, including zip code)

Registrant's telephone number, including area code: 857-246-8998

Check the appropriate box below if the Form 8-K filing is intended to simultaneously satisfy the filing obligation of the registrant under any of the following provisions:

- Written communications pursuant to Rule 425 under the Securities Act (17 CFR 230.425)
- Soliciting material pursuant to Rule 14a-12 under the Exchange Act (17 CFR 240.14a-12)
- Pre-commencement communications pursuant to Rule 14d-2(b) under the Exchange Act (17 CFR 240.14d-2(b))
- Pre-commencement communications pursuant to Rule 13e-4(c) under the Exchange Act (17 CFR 240.13e-4(c))

Indicate by check mark whether the registrant is an emerging growth company as defined in Rule 405 of the Securities Act of 1933 (17 CFR §230.405) or Rule 12b-2 of the Securities Exchange Act of 1934 (17 CFR §240.12b-2).

Emerging Growth Company

If an emerging growth company, indicate by check mark if the registrant has elected not to use the extended transition period for complying with any new or revised financial accounting standards provided pursuant to Section 13(a) of the Exchange Act.

Item 7.01 Regulation FD Disclosure.

On September 16, 2017, Pieris Pharmaceuticals, Inc. presented on the safety, tolerability and pharmacodynamics of PRS-080. The poster is furnished as Exhibit 99.1 to this Current Report on Form 8-K and is incorporated by reference herein.

The information set forth under this “Item 7.01. Regulation FD Disclosure,” including the exhibits attached hereto, shall not be deemed “filed” for purposes of Section 18 of the Securities Exchange Act of 1934, as amended, nor shall it be deemed incorporated by reference into any filing under the Securities Act of 1933, as amended, except as shall be expressly set forth by specific reference in such filing.

Item 9.01 Financial Statements and Exhibits

(d) *Exhibits.*

99.1 [Conference Poster of Pieris Pharmaceuticals, Inc., dated September 2017.](#)

SIGNATURE

Pursuant to the requirements of the Securities Exchange Act of 1934, the registrant has duly caused this report to be signed on its behalf by the undersigned hereunto duly authorized.

Dated: September 18, 2017

PIERIS PHARMACEUTICALS, INC.

By: /s/ Allan Reine

Name: Allan Reine

Title: Chief Financial Officer

EXHIBIT INDEX

Exhibit No.	Description
99.1	Conference Poster of Pieris Pharmaceuticals, Inc., dated September 2017.




Safety, tolerability, and pharmacodynamics of the hepcidin antagonist PRS-080#022-DP after single administration - a phase Ib study in anemic chronic kidney disease patients undergoing hemodialysis



Lutz Renders, MD¹, Ming Wen, MD¹, Frank Dellanna, MD¹, Heinrichs, Sven, MD¹, Klemens Budde, MD¹, Christian Rosenberger, MD¹, Christiane Erley, MD⁵, Birgit Bader, MD⁶, Claudia Sommerer, MD⁵, Schaier, Matthias, MD⁷, Werner Feuerer, MD⁸, Edgar Fenzl, MD⁹, Rachel van Swelm, PhD¹, Dorine Swinkels, MD PhD¹⁰, Klaus Kutz, MD⁹, Louis Matis, MD¹⁰, Ulrich Moebius, PhD¹¹

¹Klinikum Rechts der Isar, Department Nephrology, Munich, Germany; ²DaVita Düsseldorf, Germany; ³Charité Berlin, Germany; ⁴St. Joseph Krankenhaus, Berlin, Germany; ⁵University Hospital Heidelberg, Germany; ⁶Nuvisan Pharma Services, Neu-Ulm, Germany; ⁷FGK Clinical Research, Munich, Germany; ⁸Radboud University Medical Center, Nijmegen, The Netherlands; ⁹AccelPharm, Basel, Switzerland; ¹⁰Pieris Pharmaceuticals, Inc., Boston, Massachusetts; ¹¹Pieris Pharmaceuticals, Inc., Frising, Germany



Introduction Hepcidin plays a major role in the regulation of the iron metabolism in patients with functional iron deficiency (FID) anemia. Elevated levels of hepcidin restrict iron availability. PRS-080#022 a 20kD Anticalin® protein linked to 30kD linear poly-ethylene-glycol, is developed for the treatment of FID anemia associated with chronic kidney disease. It specifically binds to human hepcidin 25, thereby inhibiting its activity. By antagonizing hepcidin PRS-080#022 has the potential to improve iron availability and erythropoiesis, while avoiding overload with exogenous iron and reducing the administered levels of ESAs [1]. First data of this randomized, placebo controlled phase I study have already been presented on the ERA-EDTA congress in Spain, 2017 [2]. Here we show further results of single doses of PRS-080#022-DP in anemic patients with chronic kidney disease (CKD) requiring hemodialysis.

Methods and Study Design


Study design: Randomized, placebo-controlled, double-blind, multi-center study.
Main inclusion criteria: Chronic hemodialysis for ≥ 90 days, anemia of CKD, stable condition, blood hemoglobin 9.0 to 12.0 g/dL, transferrin saturation (TSAT) < 40%, ferritin > 300 ng/mL; plasma hepcidin (by mass spectrometry) 5.0 to 75 nmol/L.
Main exclusion criteria: Anemia of other cause; malignancy; infection with hepatitis B, C, or HIV; IV iron within 1 week prior to and after study medication.
Study protocol: Single IV injection of study medication; 4 weeks of follow-up; 3 cohorts with 8 patients per cohort, each cohort consisting of 6 study drug and 2 placebo treatments; increment doses of 2, 4, and 8 mg/kg from the first to the last cohort.

Treatment	N	Demographic data				Mean concentrations with standard deviation at screening		
		Mean Body weight [kg]	Mean Age [years]	Mean Hb [g/dL] [male]	Gender	Hepcidin [pmol/L]	TSAT [%]	Ferritin [ng/mL]
Placebo	6	73.32 ±10.73	54.0 ±13.2	26.50 ±4.13	3 males/3 females	31.3 ± 3.8	22.2 ±8.5	680.0 ±309.3
2 mg/kg	6	73.08 ±17.82	58.5 ±15.3	26.22 ±5.54	4 males/2 females	50.1 ±7.7	25.5 ±10.9	638.7 ±307.3
4 mg/kg	6	76.38 ±17.81	58.2 ±15.7	24.82 ±6.46	5 males/1 females	22.1 ±11.8	21.5 ±7.9	715.0 ±404.8
8 mg/kg	6	85.42 ±9.33	48.0 ±17.4	26.58 ±3.82	5 males/1 females	30.749.2	23.0 ±4.3	888.8 ±376.3
Total	24	77.08 ±14.43	54.4 ±14.1	26.56 ±4.88	17 males/7females	33.6 ±11.7	23.3 ±7.9	735.9 ±338.7

Table 1: Baseline patient characteristics

Pharmacodynamics

PRS-080#022-DP mobilizes serum iron with increases in both serum iron concentration and TSAT following treatment (Figures 1 and 2).



We have previously shown that both iron and TSAT reach maximal levels 19 hours after infusion at all 3 dose levels studied (2, 4, and 8 mg/kg) [2]. We also showed that the magnitude and duration of elevated serum iron levels and TSAT concentrations increase dose-proportionally.

As shown in Figure 1, following treatment with PRS-080#022-DP 8 mg/kg, iron levels rise and then return to baseline values between 5 and 7 days after the end of the infusion, which is a longer duration of iron mobilization when compared to the 2 and 4 mg/kg doses [2]. As also shown in Figures 1 and 2 for patients receiving 8 mg/kg PRS-080#022-DP, the TSAT and iron levels show identical time profiles, indicating that most of the iron is transferrin bound.

Compared to Placebo, Ferritin levels are not affected by the single administration of PRS-080#022-DP in all three dose groups (Figure 3). In addition, initial plasma ferritin concentrations appear to have no influence on maximal concentration of iron after the different treatments.



Figure 3: Plasma concentrations versus time profiles of FERRITIN for the three different dose levels and Placebo

Maximum serum iron mobilization appeared to correlate with baseline TSAT levels across all dose groups, such that patients with lower initial TSAT levels showed higher C_{max} iron increases following PRS-080#022-DP mediated hepcidin inhibition. The relationship between initial TSAT and C_{max} of iron is shown in Figure 4 for the 8 mg/kg dose group.




Figure 4: C_{max} of iron after infusion of 8 mg/kg PRS-080#022-DP against initial TSAT

Administration of PRS-080#022-DP resulted in a complete inhibition of free hepcidin shortly after intravenous infusion. Further, there was no evidence across all dose groups that the increase in serum iron exposure is dependent on the initial plasma hepcidin value. This is shown in Figure 5 for the 8 mg/kg dose group.




Figure 5: C_{max} of iron after infusion of 8 mg/kg PRS-080#022-DP against initial hepcidin values

Pharmacokinetics and Safety Mean plasma concentrations of Total and Free PRS-080#022-DP show a dose-dependent increase after administration of 2, 4 and 8 mg/kg PRS-080#022-DP. T_{max} of Total and Free PRS-080#022-DP occur at about 1 to 1.085 hours (median). (Figure 6)

PRS-080#022 was safe and well tolerated. The only reported serious adverse event (worsening of dry gangrene) after active treatment (2 mg/kg bw dose) was assessed as not related to PRS-080#022-DP by the investigator. No injection site reactions and no dose-dependent increase of AEs was observed within the 3 dose groups.

Conclusion The excellent safety profile and the confirmed activity of PRS-080#022-DP on iron metabolism observed in anemic dialysis dependent end-stage chronic kidney disease patients warrant further investigation of PRS-080#022-DP.




Figure 6: Mean plasma concentration of Total PRS-080#022-DP

Reference
 [1] Phase Ib Study Investigating the Safety, Tolerability, Pharmacokinetics and Pharmacodynamics Activity of the hepcidin Antagonist PRS-080#022. Results from a Randomized, Placebo Controlled, Double Blind Study Following Single Administration to Healthy Subjects. ASN 57 annual meeting & exposition, Dec. 3-6, 2015.
 [2] A phase Ib study investigating the safety, tolerability, pharmacokinetics, and pharmacodynamics of the hepcidin antagonist PRS-080#022-DP in anemic chronic kidney disease patients undergoing hemodialysis. ERA-EDTA 2017 congress Madrid, June 3-6, 2017.
 Financial disclosure statement
 L. Renders and U. Moebius are co-owner of Pieris Pharmaceuticals, Inc., all other authors have financial relationships with Pieris Pharmaceuticals, Inc. and received payment for study participation.
 Topic Chronic renal failure